



## Referenties bij MAGMA 3-2011

### Nieuwe kwaliteitsindicator heeft verkeerde noemer, p. 76

1. Button LA, Roberts SE, Evans PA, Goldacre MJ, Akbari A, Dsilva R, et al. Hospitalized incidence and case fatality for upper gastrointestinal bleeding from 1999 to 2007: a record linkage study. *Aliment Pharmacol Ther.* 2011. Jan;33(1):64-76.
2. Van Leerdam ME. Epidemiology of acute upper gastrointestinal bleeding. *Best Pract Res Clin Gastroenterol* 2008;22(2):209-24.
3. Pang SH, Ching JY, Lau JY, Sung JJ, Graham DY, Chan FK. Comparing the Blatchford and pre-endoscopic Rockall score in predicting the need for endoscopic therapy in patients with upper GI hemorrhage. *Gastrointest Endosc.* Jun;71(7):1134-40.
4. Srirajaskanthan R, Conn R, Bulwer C, Irving P. The Glasgow Blatchford scoring system enables accurate risk stratification of patients with upper gastrointestinal haemorrhage. *Int J Clin Pract.* Jun;64(7):868-74.
5. Stephens JR, Hare NC, Warshaw U, Hamad N, Fellows HJ, Pritchard C, et al. Management of minor upper gastrointestinal haemorrhage in the community using the Glasgow Blatchford Score. *Eur J Gastroenterol Hepatol.* 2009 Dec;21(12):1340-6.
6. Blatchford O, Murray WR, Blatchford M. A risk score to predict need for treatment for upper-gastrointestinal haemorrhage. *Lancet* 2000. Oct 14;356(9238):1318-21.