

DE AFDELING, ZIEKENHUIS ST. JANSDAL:
We staan als regionaal ziekenhuis dichtbij de mensen



Foto-onderschrift:

Bovenste rij (v.l.n.r.): Jaap Driest¹, Judith Amsing², Mariëlle Meijnen-Bult¹, Marleen Willems¹, Marijn Sijtsma¹, Erica Ruiter³, Daud Allajar¹.

Onderste rij: Trude Lugard-Jozuazoon⁴, Jeff de Haas⁴, Ditta Ritmeester⁴, Lena Staal-Dvaranchykava⁵, Ilona Koci-Kerkhof¹. Inzet: John Groen¹.

¹MDL-arts, ²verpleegkundig specialist MDL, ³voedingsverpleegkundige, ⁴MDL-verpleegkundige, ⁵zaalarts MDL

Referenties bij MAGMA 4-2017

TECHNIEK – Nieuwe kijk op verklevingen met CineMRI, p. 147

1. Surgical treatment of adhesion-related chronic abdominal and pelvic pain after gynaecological and general surgery: a systematic review and meta-analysis. van den Beukel BA, de Ree R, van Leuven S, Bakkum EA, Strik C, van Goor H, Ten Broek RPG. *Hum Reprod Update* 2017. May 1;23(3):276-288
2. Laparoscopic adhesiolysis in patients with chronic abdominal pain: a blinded randomised controlled multi-centre trial. Swank DJ, Swank-Bordewijk SC, Hop WC, van Erp WF, Janssen IM, Bonjer HJ, Jeekel J. *Lancet* 2003. Apr 12;361(9365):1247-51
3. Video ESSR-congres: <https://youtu.be/qi3GoaKsXuI>
4. A novel diagnostic aid for intra-abdominal adhesion detection in cine-MR imaging: Pilot study and initial diagnostic impressions. Randall D, Joosten F, ten Broek RP, Gillott R, Bardhan KD, Strik C, Prins W, van Goor H, Fenner JW. *Br J Radiol.* 2017. Jul 14:20170158. doi: 10.1259/bjr.20170158.
5. Benefits and harms of adhesion barriers for abdominal surgery: a systematic review and meta-analysis. Ten Broek RPG, Stommel MWJ, Strik C, van Laarhoven CJHM, Keus F, van Goor H. *Lancet* 2014. Jan 4;383(9911):48-59.

THEMA – Chemotherapie bij ouderen met colorectaal carcinoom, p. 150–151

1. Integraal Kankercentrum Nederland 2017. Anonymous. (Accessed 10/8, 2017, at www.cijfersoverkanker.nl).
2. Loffeld RJ, Liberov B, Dekkers PE. Yearly diagnostic yield of colonoscopy in patients age 80 years or older, with a special interest in colorectal cancer. *Geriatr Gerontol Int* 2012;12:298-303.
3. Andre T, Boni C, Mounedji-Boudiaf L, et al. Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med* 2004;350:2343-51.
4. Kuebler JP, Wieand HS, O'Connell MJ, et al. Oxaliplatin combined with weekly bolus fluorouracil and leucovorin as surgical adjuvant chemotherapy for stage II and III colon cancer: results from NSABP C-07. *J Clin Oncol* 2007;25:2198-204.
5. Haller DG, Tabernero J, Maroun J, et al. Capecitabine plus oxaliplatin compared with fluorouracil and folinic acid as adjuvant therapy for stage III colon cancer. *J Clin Oncol* 2011;29:1465-71.
6. Moertel CG, Fleming TR, Macdonald JS, et al. Levamisole and fluorouracil for adjuvant therapy of resected colon carcinoma. *N Engl J Med* 1990;322:352-8.
7. Anonymous Efficacy of adjuvant fluorouracil and folinic acid in colon cancer. International Multicentre Pooled Analysis of Colon Cancer Trials (IMPACT) investigators. *Lancet* 1995;345:939-44.
8. Sargent DJ, Goldberg RM, Jacobson SD, et al. A pooled analysis of adjuvant chemotherapy for resected colon cancer in elderly patients. *N Engl J Med* 2001;345:1091-7.
9. Wildes TM, Kallogjeri D, Powers B, et al. The Benefit of Adjuvant Chemotherapy in Elderly Patients with Stage III Colorectal Cancer is Independent of Age and Comorbidity. *J Geriatr Oncol* 2010;1:48-56.
10. Sanoff HK, Carpenter WR, Sturmer T, et al. Effect of adjuvant chemotherapy on survival of patients with stage III colon cancer diagnosed after age 75 years. *J Clin Oncol* 2012;30:2624-34.
11. Abraham A, Habermann EB, Rothenberger DA, et al. Adjuvant chemotherapy for stage III colon cancer in the oldest old: results beyond clinical guidelines. *Cancer* 2013;119:395-403.

12. McCleary NJ, Meyerhardt JA, Green E, et al. Impact of age on the efficacy of newer adjuvant therapies in patients with stage II/III colon cancer: findings from the ACCENT database. *J Clin Oncol* 2013;31:2600-6.
13. Haller DG, O'Connell MJ, Cartwright TH, et al. Impact of age and medical comorbidity on adjuvant treatment outcomes for stage III colon cancer: a pooled analysis of individual patient data from four randomized, controlled trials. *Ann Oncol* 2015;26:715-24.
14. van Eeghen EE, Bakker SD, van Bochove A, Loffeld RJ. High Risk Stage 2 and Stage 3 Colon Cancer, Predictors of Recurrence and Effect of Adjuvant Therapy in a Nonselected Population. *Int Sch Res Notices* 2015;2015:790186.
15. Andre T, Quinaux E, Louvet C, et al. Phase III study comparing a semimonthly with a monthly regimen of fluorouracil and leucovorin as adjuvant treatment for stage II and III colon cancer patients: final results of GERCOR C96.1. *J Clin Oncol* 2007;25:3732-8.
16. Shi Q, Sobrero AF, Shields AF, et al. Prospective pooled analysis of six phase III trials investigating duration of adjuvant oxaliplatin-based therapy (3 vs 6 months) for patients with stage III colon cancer: The IDEA (International Duration Evaluation of Adjuvant chemotherapy) collaboration. *Abstract presented at ASCO 2017*.
17. Venderbosch S, Doornbehal J, Teerenstra S, Lemmens W, Punt CJ, Koopman M. Outcome of first line systemic treatment in elderly compared to younger patients with metastatic colorectal cancer: a retrospective analysis of the CAIRO and CAIRO2 studies of the Dutch Colorectal Cancer Group (DCCG). *Acta Oncol* 2012;51:831-9.
18. van Steenbergen LN, Lemmens VE, Rutten HJ, Wymenga AN, Nortier JW, Janssen-Heijnen ML. Increased adjuvant treatment and improved survival in elderly stage III colon cancer patients in The Netherlands. *Ann Oncol* 2012;23:2805-11.

THEMA – IJzergebreksanemie. Wat te doen bij kwetsbare ouderen, p. 153

1. Joosten E, Pelemans W, Hiele M et al. Prevalence and causes of anaemia in a geriatric hospitalized population. *Gerontology* 1992;38:111-117.
2. Hamaker ME, Acampo T, Remijn JA et al. Diagnostic choices and clinical outcomes in octogenarians and nonagenarians with iron-deficiency anemia in the Netherlands. *J Am Geriatr Soc* 2013;61:495-501.
3. Luman W, Ng KL. Audit of investigations in patients with iron deficiency anaemia. *Singapore Med J* 2003;44:504-510.
4. Lindsay JO, Robinson SD, Jackson JE et al. The investigation of iron deficiency anemia--a hospital based audit. *Hepatogastroenterology* 1999;46:2887-2890.
5. Stephens MR, Hopper AN, White SR et al. Colonoscopy first for iron-deficiency anaemia: a Numbers Needed to Investigate approach. *QJM* 2006;99:389-395.
6. Majid S, Salih M, Wasaya R et al. Predictors of gastrointestinal lesions on endoscopy in iron deficiency anemia without gastrointestinal symptoms. *BMC Gastroenterol* 2008;8:52.
7. Chatrenet P, Friocourt P, Ramain JP et al. Colonoscopy in the elderly: a study of 200 cases. *Eur J Med* 1993;2:411-413.
8. Lukens FJ, Loeb DS, Machicao VI et al. Colonoscopy in octogenarians: a prospective outpatient study. *Am J Gastroenterol* 2002;97:1722-1725.
9. Avgerinos DV, Llaguna OH, Lo AY et al. Evolving management of colonoscopic perforations. *J Gastrointest Surg* 2008;12:1783-1789.
10. Jafri SM, Arora A. Silent perforation: an iatrogenic complication of colonoscopy. *Surg Laparosc Endosc Percutan Tech* 2007;17:452-454.

11. Jafri SM, Monkemuller K, Lukens FJ. Endoscopy in the elderly: a review of the efficacy and safety of colonoscopy, esophagogastroduodenoscopy, and endoscopic retrograde cholangiopancreatography. *J Clin Gastroenterol* 2010;44:161-166.
12. Jonker JM, Hamaker ME, Soesan M et al. Colon cancer treatment and adherence to national guidelines: does age still matter? *J Geriatr Oncol* 2012;3:131-137.
13. Gooiker GA, Dekker JW, Bastiaannet E et al. Risk Factors for Excess Mortality in the First Year After Curative Surgery for Colorectal Cancer. *Ann Surg Oncol* 2012.
14. Cull JD, Kukreja S, Hieken TJ. Surgical treatment of colorectal cancer in the advanced elderly: does outcome justify treatment? *J.Clin.Oncol. ASCO meeting abstracts*, 3629. 9-6-2012.
15. Ronning, Wyller TB, Jordhoy MS et al. Frailty indicators and functional status in older patients after colorectal cancer surgery. *J Geriatr Oncol* 2014;5:26-32.

THEMA – Oog voor oudere patiënt voor, tijdens en na multidisciplinair overleg, p. 154–157

1. R. O’Caoimh et al. Advance care planning within survivorship care plans for older cancer survivors: A systematic review. 23-06-2017. *Pubmed*. <https://www.ncbi.nlm.nih.gov/pubmed/28687135>
2. E.J. Guerard et al. Survivorship care for older adults with cancer: U13 conference report. *J Geriatr Oncol.* 2016 Jul;7(4):305-12.
3. Cath Taylor, Jennifer Finnigan-John and James Green. "No decision about me without me" in the context of cancer multidisciplinary team meetings: a qualitative interview study. *BMC Health Serv Res* 2014; 14:488-462.
4. Josep M. Barras et al. Policy statement on multidisciplinary cancer care. European partnership action against cancer consensus group. *European Journal of Cancer* 2014; Feb;50(3):475-80.
5. Pype P, Mertens F, Belche J et al. Experiences of hospital-based-multidisciplinary team meetings in oncology: An interview study among participating general practitioners.

THEMA – Prehabilitatie voor oudere darmkankerpatiënt, p. 158–159

1. Topp R, Ditmyer M, King K, Doherty K, Hornyak J, 3rd. The effect of bed rest and potential of prehabilitation on patients in the intensive care unit. *AACN Clin Issues* 2002;13(2):263-76.
2. Ditmyer MM, Topp R, Pifer M. Prehabilitation in preparation for orthopaedic surgery. *Orthop Nurs.* 2002;21(5):43-51; quiz 2-4.
3. Hulzebos EH, Smit Y, Helders PP, van Meeteren NL. Preoperative physical therapy for elective cardiac surgery patients. *The Cochrane database of systematic reviews* 2012; 11:Cd010118.
4. Robinson TN, Wu DS, Pointer LF, Dunn CL, Moss M. Preoperative cognitive dysfunction is related to adverse postoperative outcomes in the elderly. *J Am Coll Surg.* 2012;215(1):12-7; discussion 7-8.
5. Gooiker GA, Dekker JW, Bastiaannet E, van der Geest LG, Merkus JW, van de Velde CJ, et al. Risk factors for excess mortality in the first year after curative surgery for colorectal cancer. *Ann Surg Oncol.* 2012;19(8):2428-34.
6. Robinson TN, Wu DS, Sauaia A, Dunn CL, Stevens-Lapsley JE, Moss M, et al. Slower walking speed forecasts increased postoperative morbidity and 1-year mortality across surgical specialties. *Ann Surg.* 2013;258(4):582-8; discussion 8-90.

7. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987;40(5):373-83.
8. Snowden CP, Prentis J, Jacques B, Anderson H, Manas D, Jones D, et al. Cardiorespiratory fitness predicts mortality and hospital length of stay after major elective surgery in older people. *Ann Surg.* 2013;257(6):999-1004.
9. Heldens A, Bongers BC, Lenssen AF, Stassen LPS, Buhre WF, van Meeteren NLU. The association between performance parameters of physical fitness and postoperative outcomes in patients undergoing colorectal surgery: An evaluation of care data. *Eur J Surg Oncol.* 2017.
10. Sui X, LaMonte MJ, Laditka JN, Hardin JW, Chase N, Hooker SP, et al. Cardiorespiratory fitness and adiposity as mortality predictors in older adults. *JAMA* 2007;298(21):2507-16.
11. Dekker JW, van den Broek CB, Bastiaannet E, van de Geest LG, Tollenaar RA, Liefers GJ. Importance of the first postoperative year in the prognosis of elderly colorectal cancer patients. *Ann Surg Oncol.* 2011;18(6):1533-9.
12. Ren L, Zhu D, Wei Y, Pan X, Liang L, Xu J, et al. Enhanced Recovery After Surgery (ERAS) program attenuates stress and accelerates recovery in patients after radical resection for colorectal cancer: a prospective randomized controlled trial. *World J Surg.* 2012;36(2):407-14.
13. Mosquera C, Spaniolas K, Fitzgerald TL. Impact of frailty on approach to colonic resection: Laparoscopy vs open surgery. *World J Gastroenterol.* 2016;22(43):9544-53.
14. Cerdan Santacruz C, Frasson M, Flor-Lorente B, Ramos Rodriguez JL, Trallero Anoro M, Millan Scheiding M, et al. Laparoscopy may decrease morbidity and length of stay after elective colon cancer resection, especially in frail patients: results from an observational real-life study. *Surg Endosc.* 2017.
15. Gietelink L, Wouters MW, Bemelman WA, Dekker JW, Tollenaar RA, Tanis PJ. Reduced 30-Day Mortality After Laparoscopic Colorectal Cancer Surgery: A Population Based Study From the Dutch Surgical Colorectal Audit (DSCA). *Ann Surg.* 2016;264(1):135-40.
16. Bruns ER, van den Heuvel B, Buskens CJ, van Duijvendijk P, Festen S, Wassenaar EB, et al. The effects of physical prehabilitation in elderly patients undergoing colorectal surgery: a systematic review. *Colorectal Dis.* 2016.
17. O'Doherty AF, West M, Jack S, Grocott MP. Preoperative aerobic exercise training in elective intra-cavity surgery: a systematic review. *Br J Anaesth.* 2013;110(5):679-89.
18. Hijazi Y, Gondal U, Aziz O. A systematic review of prehabilitation programs in abdominal cancer surgery. *Int J Surg.* 2017;39:156-62.
19. Moran J, Guinan E, McCormick P, Larkin J, Mockler D, Hussey J, et al. The ability of prehabilitation to influence postoperative outcome after intra-abdominal operation: A systematic review and meta-analysis. *Surgery.* 2016;160(5):1189-201.
20. Barberan-Garcia A, Ubre M, Roca J, Lacy AM, Burgos F, Risco R, et al. Personalised Prehabilitation in High-risk Patients Undergoing Elective Major Abdominal Surgery: A Randomized Blinded Controlled Trial. *Ann Surg.* 2017.
21. ESSO/SIOG. GO SAFE (Geriatric Oncology Surgical Assessment and Functional rEcovery after Surgery).
Meer informatie op <http://wwwessoneweborg/news/gosafe-project/>.
22. Academy L. Life and Vitality Assessment (LAVA). Beschikbaar via:
<http://wwwleydenacademynl/life-and-vitality-assessment/>.
European J of Gen Practice 2017; Dec: 23(1) 155-163.

THEMA – Hoe blijf je als arts vitaal? p. 159–161

1. <https://www.cbs.nl/nl-nl/achtergrond/2017/14/beroep-en-werkdruk>
2. <https://www.cbs.nl/nl-nl/achtergrond/2017/14/beroep-en-werkdruk>
3. Wallace, J.E., Lemaire, J.B., & Ghali, W.A. Physician wellness: a missing quality indicator. *The Lancet* 2009, 374 (9702), 1714-1721.
4. Shanafelt, T.D., Balch, C. M., Bechamps, G., Russell, T., Dyrbye, L., Satele, D., ... & Freischlag, J. Burnout and medical errors among American surgeons. *Annals of surgery* 2010, 251(6), 995-1000.
5. Halbesleben, J.R., & Rathert, C. Linking physician burnout and patient outcomes: exploring the dyadic relationship between physicians and patients. *Health care management review* 2008, 33(1), 29-39.
6. Shanafelt, T.D., Hasan, O., Dyrbye, L.N., Sinsky, C., Satele, D., Sloan, J., & West, C.P. Changes in burnout and satisfaction with work-life balance in physicians and the general US working population between 2011 and 2014. *Mayo Clinic Proceedings* 2015 (December, Vol. 90, No. 12, pp. 1600-1613). Elsevier.
7. <https://www.medischcontact.nl/nieuws/laatste-nieuws/artikel/arbeidsbelasting-artsen-dubbel-zo-groot.htm>
8. <https://www.medischcontact.nl/nieuws/laatste-nieuws/artikel/oudere-specialisten-maken-geen-haast-met-hun-pensioen.htm>
9. [https://www\(skb.nl/nl/nieuws/oudere-werknemer-verrassend-vitaal?utm_medium=email&utm_campaign=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal&utm_content=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal+CID_db44ee999c0efa1ac26382364c370c6f&utm_source=Mailing%20%20More&utm_term=vitaliteit%20ouderen%20vs%20jongeren](https://www(skb.nl/nl/nieuws/oudere-werknemer-verrassend-vitaal?utm_medium=email&utm_campaign=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal&utm_content=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal+CID_db44ee999c0efa1ac26382364c370c6f&utm_source=Mailing%20%20More&utm_term=vitaliteit%20ouderen%20vs%20jongeren)
10. Bakker, A.B., & Demerouti, E. The job demands-resources model: State of the art. *Journal of managerial psychology* 2007, 22(3), 309-328.
11. West C.P., Dyrbye L.N., Erwin P.J., Shanafelt T.D.. Interventions to prevent and reduce physician burnout: a systematic review and meta-analysis. *The Lancet* 2016; 388(10057):2272-2281.
12. Panagioti M, Panagopoulou E, Bower P, et al. Controlled Interventions to Reduce Burnout in Physicians: A Systematic Review and Meta-analysis. *JAMA Internal Medicine* 2016.
13. [https://www\(skb.nl/nl/nieuws/oudere-werknemer-verrassend-vitaal?utm_medium=email&utm_campaign=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal&utm_content=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal+CID_db44ee999c0efa1ac26382364c370c6f&utm_source=Mailing%20%20More](https://www(skb.nl/nl/nieuws/oudere-werknemer-verrassend-vitaal?utm_medium=email&utm_campaign=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal&utm_content=SKB%20Nieuwsbrief%20-%20Oudere%20werknemer%20verrassend%20vitaal+CID_db44ee999c0efa1ac26382364c370c6f&utm_source=Mailing%20%20More)

LEVER – Leververvetting: geen onschuldige aandoening, p. 165–169

1. <https://www.kennisinstituut.nl/sites/default/files/NVMDL%20Kennisagenda.pdf>
2. Data van CBS StatLine, the Netherlands.
3. volksgezondheidzorg.info 2015.
4. Gezondheidsenquete/leefstijlmonitor CBS i.s.m. RIVM en Trimbos-instituut 2016.
5. van Meer S, van Erpecum KJ, Sprengers D, Klümpen HJ, Jansen PL, Ijzermans JN, Siersema PD, de Man RA, Verheij J. Hepatocellular carcinoma in noncirrhotic livers is associated with steatosis rather than steatohepatitis: potential implications for pathogenesis. *Eur J Gastroenterol Hepatol* 2016 Aug;28(8):955-62.

6. Koek GH, Liedorp PR, Bast A. The role of oxidative stress in non-alcoholic steatohepatitis. *Clin Chim Acta* 2011;412:1297-1305.
7. Angulo P, Bugianesi E, Bjornsson ES, Charatcharoenwithaya P, Mills PR, Barrera F, Haflidadottir S, Day CP, George J. Simple noninvasive systems predict long-term outcomes of patients with nonalcoholic fatty liver disease. *Gastroenterology* 2013 Oct;145(4):782-9.
8. Verhaegh P, Bavia R, Winkens B, Mascllee A, Jonkers D, Koek G. Non-invasive Tests Do Not Accurately Differentiate Non-alcoholic Steatohepatitis from Simple Steatosis: A Systematic Review and Meta-Analysis. *Clin Gastroenterol Hepatol* 2017; Aug: S1542-3565.
9. Vilar-Gomez E, Martinez-Perez Y, Calzadilla-Bertot L, et al. Weight loss through lifestyle modification significantly reduces features of nonalcoholic steatohepatitis. *Gastroenterology* 2015;149:367–378. e365; quiz e314–365.
10. Neuschwander-Tetri BA, Loomba R, Sanyal AJ, et al. Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet* 2015;385:956–965.
11. Cariou B, Hanf R, Lambert-Porcheron S, et al. Dual peroxisome proliferator-activated receptor alpha/delta agonist GFT505 improves hepatic and peripheral insulin sensitivity in abdominally obese subjects. *Diabetes Care* 2013;36:2923–2930.

CASUÏSTIEK – Ischemie van de maag, p. 171

1. Tang SJ, Daram SR, WuR, Bhajee F. Pathogenesis, diagnosis, and management of gastric ischemia. *Clin gastroenterol Hepatol* 2014. Feb; 12(2), 246-52
2. Lunca S, Rikkers A, Stanescu A. Acute massive gastric dilatation: severe ischemia and gastric necrosis without perforation. *Rom J Gastroenterol* 2005. Sep ; 14(3): 279-83.
3. Mishima T, KoharabN,Tajima Y, Maeda J, Inoue K, Ohno T, Kitasato A, Watanabe T,Irie J, Adachi T, Kuroki T, Eguchi S, Kanematsu T. Gastric rupture with necrosis following acute gastric dilatation: report of a case. *Surg Today* 2012. Oct;42(10):997-1000
4. Kolkman JJ, Reeders JW AJ. Gastro-intestinale chirurgie en gastro-enterologie. VIII. Gastro-enterologische aspecten van chronische maag-darmischemie. *Ned Tijdschr Geneesk*. 2000;144:813-4.